



PHYSICAL CHEMISTRY 2016

*13th International Conference on
Fundamental and Applied Aspects of
Physical Chemistry*

*Proceedings
Volume I*

BELGRADE
September 26 - 30, 2016

ISBN 978-86-82475-34-7

Title: Physical Chemistry 2016 (Proceedings)

Editors: Željko Čupić and Slobodan Anić

Published by: Society of Physical Chemists of Serbia, Studentski trg 12-16, 11158, Belgrade, Serbia.

Publisher: Society of Physical Chemists of Serbia

For Publisher: S. Anić, President of Society of Physical Chemists of Serbia

Printed by: "Jovan", Printing and Publishing Company; 200 Copies.

Number of pages: 6+502; Format B5; printing finished in September 2016

Text and Layout: "Jovan"

Neither this book nor any part may be reproduced or transmitted in any form or by any means, including photocopying, or by any information storage and retrieval system, without permission in writing from the publisher.

200 - Copy printing



PHYSICAL CHEMISTRY 2016

*13th International Conference on
Fundamental and Applied Aspects of
Physical Chemistry*

Organized by

*The Society of Physical Chemists of
Serbia*

in co-operation with

Institute of Catalysis Bulgarian Academy of Sciences

and

*Boriskov Institute of Catalysis Siberian Branch of
Russian Academy of Sciences*

and

University of Belgrade, Serbia:

*Faculty of Physical Chemistry
Institute of Chemistry, Technology and Metallurgy
Vinča Institute of Nuclear Sciences
Faculty of Pharmacy
Institute of General and Physical Chemistry, Belgrade, Serbia*

International Organizing Committee

Chairman: S. Anić (Serbia)
Vice-chairman: M. Gabrovska (Bulgaria)
A. A. Vedyagin (Russia)
S. N. Blagojević (Serbia)
Members: N. Cvjetičanin (Serbia), S. M. Blagojević (Serbia), M. Daković (Serbia), J. Dimitrić-Marković (Serbia), T. Grozdić (Serbia), Lj. Ignjatović (Serbia), D. Jovanović (Serbia), J. Jovanović (Serbia), M. Kuzmanović (Serbia), D. Marković (Serbia), B. Milosavljević (USA), M. Mojović (Serbia), N. Ostrovski (Serbia), N. Pejić (Serbia), M. Petković (Serbia), A. Popović-Bjelić (Serbia), B. Simonović (Serbia), D. Stanisavljev (Serbia), M. Stanković (Serbia), Z. Šaponjić (Serbia), B. Šljukić (Serbia), G. Tasić (Serbia), N. Vukelić (Serbia), V. Vukojević (Sweden)

International Scientific Committee

Chairman: Ž. Čupić (Serbia)
Vice-chairmans: V. N. Parmon (Russia)
S. Rakovsky (Bulgaria)
B. Adnađević (Serbia)
Members: S. Anić (Serbia), A. Antić-Jovanović (Serbia), G. Bačić (Serbia), R. Cervellati (Italy), G. Ćirić-Marjanović (Serbia), A. Cricenti (Italy), V. Dondur (Serbia), S. D. Furrow (USA), L. Gábor (Hungary), Vilmos Gáspár (Hungary), K. Hedrih (Serbia), M. Jeremić (Serbia), E. Kiš (Serbia), Lj. Kolar-Anić (Serbia), U. Kortz (Germany), T. Kowalska (Poland), V. Kuntić (Serbia), Z. Marković (Serbia), S. Mentus (Serbia), K. Novaković (UK), B. Novakovski (Poland), T. Parac Vogt (Belgium), M. Perić (Serbia), M. Plavšić (Serbia), G. Schmitz (Belgium), I. Schreiber (Czech Republic), P. Ševčík (Slovakia), N. Stepanov (Russia), M. Trtica (Serbia), V. Vasić (Serbia), D. Veselinović (Serbia), Á. Tóth (Hungary)

Local Executive Committee

Chairman: S. N. Blagojević
Vice-chairmans: A. Ivanović-Šašić
A. Stojiljković
Members: M. Ajduković, P. Banković, N. Bošnjaković, I. N. Bujanja, D. Dimić, A. Dobrota, J. Dostanić, A. Ignjatović, S. Jovanović, Z. Jovanović, A. Jović, N. Jović-Jovičić, D. Lončarević, M. Kragović, J. Krstić, S. Maćešić, J. Maksimović, V. Marković, D. Milenković, M. Milovanović, B. Nedić-Vasiljević, M. Pagnacco, A. Pavićević, N. Potkonjak, D. Ranković, M. Ristić, B. Stanković, A. Stanojević

ON THE USE OF VOLTAMMETRIC DATA IN BIFURCATION ANALYSIS OF ELECTROCHEMICAL OSCILLATORY SYSTEM: Cu | 1.0 M TFA

N. I. Potkonjak¹, T. N. Potkonjak² and Z. Nikolić³

¹ *Chemical Dynamics Laboratory, Vinča Institute of Nuclear Sciences,
University of Belgrade, P.O. Box 522, 11001 Belgrade, Serbia.
(npotkonjak@vinca.rs)*

² *Urban Planning Institute of Belgrade, 30 Palmotićeve, 11000 Belgrade,
Serbia*

³ *Faculty of Physics, 12 Studentski trg, 11000 Belgrade, Serbia*

ABSTRACT

Bifurcation analysis of the Cu | 1 M TFA electrochemical oscillatory system was done by using voltammetric data, obtained under quasi-potentiostatic polarization conditions. A super-critical Hopf bifurcation and a saddle-loop bifurcation were identified at following bifurcation potentials $E_{\text{BIF1}} = 0.5446$ V and $E_{\text{BIF2}} = 0.7536$ V, respectively.

INTRODUCTION

Being far-from-equilibrium, the electrochemical systems can spontaneously form a wide spectrum of spatio-temporal patterns [1, 2]. Spontaneous emergence of such patterns implies that the dynamical systems, such as electrochemical, undergo self-organization [2]. Appearance of oscillations implies that the electrochemical system can no longer achieve stable steady states, and that its existence is only possible through oscillatory states [1]. Transient from a stable steady state to an oscillatory state goes via bifurcation point. In order to be identified, evolution of the system has to be monitored through phase space diagram; constructed from a series of coordinates characterized by a parameter (controlled quantity) and a variable (quantity which is an appropriate representative of the system's state) [3,4]. Looking from this perspective, linear sweep voltammetry can be observed as instrumental technique which dynamically drives the system throughout a series of variable-to-parameter points [5]. Therefore, voltammogram is in fact a phase space diagram. In this study, a copper electrode in a trifluoroacetic acid (TFA) will be anodically polarized by means of linear sweep voltammetry, under quasi-potentiostatic polarization conditions. Bifurcation analysis will be conducted by using voltammetric data.

EXPERIMENTAL

Experiments were carried out in a three-electrode electrolytic cell, at 293 K, with a copper rod (Goodfellow, 99.99%) as the working electrode, Pt foil as the counter electrode, and a saturated calomel electrode (SCE) as the reference one. The working electrode was embedded in a plastic capillary, leaving only the rod cross-section (0.0314 cm^2) exposed to the electrolyte solution. Prior to each experiment the working electrode was abraded by a series of wet sanding paper with different grit size (320, 600, 800, 1000, 1200 and 2000). Thereupon, the working electrode was rinsed with deionized water in an ultrasonic bath for 2 min. A Luggin capillary was used. Electrolyte solution was 1 M TFA. Experiments were carried under natural convection of electrolyte in the electrolytic cell, without any external resistance applied in the circuit. Linear sweep voltammetry was performed using anodic scan, starting from $0.5 \text{ V}_{\text{SCE}}$, at a rate of 1 mV s^{-1} . All applied circuit potentials (E) are given with the respect to the SCE.

RESULTS AND DISCUSSION

By carefully selection of initial conditions, we were able to locate an oscillatory state (OsS) region of investigated system at the current-potential (I - E) polarization curve, Fig.1. The first bifurcation point occurs at $E_{\text{BIF1}} = 0.5446 \text{ V}$ (looking from the direction of polarization). This bifurcation point can be observed as an entering one, marking a discontinuity in evolution of the system from the steady stable state (SSS1) to OsS; it is a characteristic of the $\text{SSS1} \rightarrow \text{OsS}$ transient.




Figure 1. I - E polarization curve of Cu electrode in 1M TFA.

With respect to above mention perspective, the second bifurcation point, located at $E_{\text{BIF2}} = 0.7536 \text{ V}_{\text{SCE}}$, can be taken as the exit point; it is a feature of the $\text{OsS} \rightarrow \text{SSS2}$ transient, Fig.1.

Smooth increase of current oscillation amplitudes observed near on the $\text{SSS1} \rightarrow \text{OsS}$ transient is shown in Fig.2 (a). Linear dependence between the square amplitude of current oscillations (A^2) and increase of the distance of controllable parameter from E_{BIF1} ($|E_{\text{OSC}} - E_{\text{BIF1}}|$) was found, Fig.2 (b). Along with the lack of hysteresis at the $\text{SSS1} \rightarrow \text{OsS}$ transient[5] (not shown in abstract), suggests on super-critical Hopf (SUPH) bifurcation[5, 6].




Figure 2. (a) Detail from I - E polarization curve, Fig.1, in the vicinity of the $\text{SSS1} \rightarrow \text{OsS}$ transient. (b) Dependence of the square amplitude of current oscillations (A^2) vs. distance of controllable parameter from E_{BIF1} ($|E_{\text{OSC}} - E_{\text{BIF1}}|$). (c) Detail from I - E polarization curve, Fig.1, in the vicinity of the $\text{OsS} \rightarrow \text{SSS2}$ transient (units of x-axis are converted from potential to time). (d) Semilogarithmic plot of the period current oscillations (τ) vs. the distance of controllable parameter from E_{BIF2} ($\ln |E_{\text{OSC}} - E_{\text{BIF2}}|$).

On the other hand, the period of current oscillations (τ) increases as the system approaches the $\text{OsS} \rightarrow \text{SSS2}$ transient, ending with sudden termination of oscillatory behaviour, Fig.2 (c). Linear dependence between τ

and logarithmic value of the distance from E_{BIF2} ($\ln |E_{OSC} - E_{BIF2}|$) is shown in Fig. 2 (d). Combined with an existence of hysteresis [5] (not shown in abstract), it implies that the OsS \rightarrow SSS2 transient goes via the saddle-loop bifurcation [3, 5, 6]. Correlation coefficients (R), shown in Fig. 2(b) and 2(d), were found to be considerably high. Hence, we can say that the procedure for bifurcation analysis was verified.

CONCLUSION

In this study, a voltammetric data, obtained from quasi-potentiostatic polarization curve of the Cu | 1M TFA electrochemical oscillatory system, were used for bifurcation analysis. This was based on the presumption that voltammogram can be considered as a phase space diagram. Results presented have showed that bifurcation scenario can be built from voltammetric data with a high accuracy.

Acknowledgement

This work was supported by the Ministry of Education, Science and Technological Development of Republic of Serbia (Grants no. 172015).

REFERENCES

- [1] M. Orlik, Self-Organization in electrochemical systems I: General principles of self-organization. Temporal Instabilities, Springer-Verlag, Berlin, Heidelberg, 2012
- [2] K. Krischer, N. Mazouz, P. Grauel, *Angewandte Chemie*, 2001, **40**, 850-869.
- [3] M.T.M. Koper, *Journal of Chemical Society, Faraday Transactions*, 1998, **94**, 1369-1378.
- [4] D. Kondepudi, I. Prigogine, *Modern Thermodynamics: from heat engines to dissipative structures*, John Wiley & Sons Ltd, Baffins Lane, Chichester, West Sussex PO19 1UD, England, 1998.
- [5] N. Potkonjak, Ph.D. Thesis, Faculty of Physical Chemistry, University of Belgrade, Belgrade, 2015 (in Serbian).
- [6] D. Sazou, M. Pagitsas, *Journal of Electroanalytical Chemistry* 1998, **451**, 77-87.

544(082)

66.017/.018(082)

502/504(082)

663/664:658.56(082)

615.31:547(082)

INTERNATIONAL Conference on Fundamental and Applied Aspects of Physical Chemistry (13 ; 2016 ; Beograd)

Physical Chemistry 2016 : proceedings. Vol. 2 / 13th International Conference on Fundamental and Applied Aspects of Physical Chemistry, Belgrade, 26-30 September 2016 ; [editors Željko Čupić and Slobodan Anić]. - Belgrade : Society of Physical Chemists of Serbia, 2016 (Belgrade : Jovan). - IV, 507-930 str. : ilustr. ; 24 cm

Tiraž 200. - Bibliografija uz svaki rad.

ISBN 978-86-82475-33-0

1. Society of Physical Chemists of Serbia (Beograd)

а) Физичка хемија - Зборници б) Наука о материјалима - Зборници с) Животна средина - Заштита - Зборници д) Животне намирнице - Контрола квалитета - Зборници е) Фармацеутска хемија - Зборници
COBISS.SR-ID 225802508

08.09.2016

